# Appendix 17 Soil Sampling Procedures

- 1. The area to be sampled should be divided into obviously different soil and site types and each type sampled as a separate unit. A 6 " OS map must show the total area under proposal, show the pattern of sampling and make clear what sections are represented by what samples.
- 2. For each unit, if there are clearly different horizons then sample each horizon separately. Otherwise take separate samples from the layers 0-20 cm (surface) and 20 40 cm (subsurface). In the absence of clearly defined soil horizons it is possible to manipulate the sampling procedure and analysis results by taking the second sample as near as possible to the 20 cm point. This difficulty can be overcome by taking an additional sample from the zone below 40 cm. This will involve a third sample for that particular site but will also remove any doubts about the result of the soil analysis.
- 3. From each horizon or layer collect 12 15 or more samples. Each of these is treated as a subsample.
- 4. Collect the subsamples by travelling across the area to be sampled using a zigzag or "S" shaped route. All sampling locations should be identified on 6" OS map, if available.
- 5. Combine the 12 to 15+ sub samples from each horizon or layer in a large clean bag and mix thoroughly. Make sure to keep the horizon or layer sample separate and clearly labelled.
- 6. Take a portion of the mixed sub-samples, at least 100 g (40zs) and send to a laboratory with identifying and other details.
- 7. Record all aspects of sampling procedure and site details. The location of each sample unit and the location of where the sub–sample were taken should be recorded on a 6 " OS map.
- 8. The samples should be assessed in the laboratory for pH and free CaCO<sub>3</sub>.
- 9. Record the depth of any apparent calcareous layer in the soil profile.
- 10. Sampling should be done by an approved forester. These foresters should have acquainted themselves fully with the recognised and accepted forestry soil sampling techniques as published by Forest Service.
- 11. The relevant Forest Service Inspector should receive a copy of the soil sampling form fully completed and the soil sampling map showing the sampling areas or units. Each sample must be related to a plot number and each zone of sampling clearly identified. S/he should also receive a copy of the laboratory soil analysis and comments including species recommendations, site suitability etc. The completed soil sampling form, species recommendations, together with the map and drainage report if applicable, must be signed by the soil scientist making the recommendation
- 12. The procedures as outlined above do not cover man made soils e.g. land fill or where there has been extensive reclamation, levelling, drainage or disturbance of some kind. Where there has been interference by man the soil sampler should discuss the soil sampling procedure with the relevant soil analysis laboratory before any samples are taken.

#### **Explanatory Notes**

If the crop fails or does not perform satisfactorily and this performance is shown to be related to edaphic conditions then the soil sampling procedures and analysis will be questioned. The responsibility rests with the land owner, the site developer and the soil sample laboratory. Because of the possibility of crop failure it is recommended that an independent third party should take the soil samples.

The Forest Service at this stage will conduct its own investigation including soil sampling and independent analysis. If these analysis results are at variance with the first lot the land owner will be advised of this fact. Forest Service will issue a written warning to the soil sampler at this stage. If in the event of a repeat happening the soil sampler will be told that s/he will no longer be considered a suitable and fit person to submit samples and reports.

Following on from here, if in the opinion of the Forest Service the land does not have the potential to produce a crop of minimum YC 14 Sitka spruce grant money and premiums will have to be refunded to the Department. Furthermore the land owner will be very likely to seek adequate compensation for his/her loss of income while the land was out of agricultural production.

Two forest soil laboratories offer a soil analysis service. They are:

Coillte Teo U.C.D

Site Studies Section Dept of Environmental Resource Management

Coillte Research Laboratory University College

Newtownmountkennedy Belfield Co Wicklow Dublin 4

Tel: 01-2819493 Tel: 01-7067716

# **SOIL SAMPLING FORM**

## PERSONAL DETAILS

Name & Address			
Telephone No.	Fax. No.	Mobile No.	
Your Reference			
Sampled by		Date Sampled	
Townland		OS Grid Ref.	
SITE DETAILS			
Is it currently planted?	Year planted?	What species?	
Area: (Ha)	(Acre)	Dominant ground vegetation?	
Aspect: (facing N,S,E,W, or flat)		Elevation: (1) ft or (2) m	
PREVIOUS LAND USE	1	1	
Farming Practices before			
Previous Crop			
Fertilised/limed Yes/No	Year	Type/Rate	
Relcaimed Yes/No	Year	Туре	
SOIL SAMPLING DEPTHS			
Topsoil depth: (cm)	SUBSOIL DEPTH		
Situation 1. Mineral (depth to Calcareous material	(cm))		
12345678	910111213	_141516	
Situation 2. Peat over Calcareous material Depth to	Calcareous material (cm)		
12_3_4_5_6_7_8	910111213	141516	
ANY OTHER COMMENTS			

### SITE DETAILS: (TICK AS APPROPRIATE)

TOPOGRAPHY Flat Concave Convex Bottom-slope Mid-slope Top-slope	SOIL TYPE Brown Earth Brown Podsolic Podsol Podsol+pan Lithosol Peaty Gley Gley Blanket Bog Raised Bog Fen peat over calcareous	CALCAEROUS MAT Shell Marl Marl (shells) Till Coarse (+boulders) Fine (-boulders) Esker	TERIAL	Mod. exposed  Mod. sheltered		
SITE FERTILITY* Class A Class B Class C Class X		Potential Poor Mod. Good No	SITI Floo Frost			
*EXPLANATION OF SITE FERTILITY  Class A: *Fields and ornamental ground*. These are areas, which have been in intensive agricultural use up to relatively recent times, so that they carry characteristic agricultural vegetation (pasture grasses and herbaceous plants, often with high proportion of rushes. They are among the most fertile site types.  Class B: *Furze or whin*. These are sites that were once enclosed by banks, walls or ditches. This indicates that at one time they were considered sufficiently fertile to justify bringing them under agricultural use, and were probably cultivated. The class would include all long-abandoned agricultural land (indicated by the presence of Ulex or bracken). It might also be extended, on the baiss of local knowledge, and experience, to include unenclosed areas on mineral soils derived from parent materials of shale, mica-schist or gramitic origin.  Class C:*Rough pasture, with or without cropping rock*. These are areas of unenclosed ground, which have never been cultivated or brought under any form of intensive agricultural use. Sites to include are those on unenclosed land, usually upland or bogland, carrying typical unimproved heath land or peat land vegetation.  Class X:*Woodland*. Sites to be included are coniferous, broadleaved and mixed woodland.						
LABORATORY TESTS: TICK THE TEST THAT YOU REQUIRE ON YOUR SOIL SAMPLE  1. pH, e.g. for the purpose of determining site suitability for forestry 2. pH & Calcium, e.g. for the purpose of determining site suitability & species selection						
3. pH, Phosphorous, Potassium, Calcium & Magnesium, e.g. for the purpose of determining growth problems and preparing fertiliser prescriptions						
*Please note that a "topsoil" a SEND SAMPLES TO:	Site Studies Labora Newtownmountke Co. Wicklow. Tel: 01-2819493,	ratory, ennedy, Fax: 01-2810465 mental Resource Manage	ment,			